

Noncommutativity in a Time-Dependent Background

Louise Dolan¹

Department of Physics

University of North Carolina, Chapel Hill, NC 27599-3255

Chiara R. Nappi²

Department of Physics, Jadwin Hall

Princeton University, Princeton, NJ 08544-0708

We compute a time-dependent noncommutativity parameter in a model with a time-dependent background, a space-time metric of the plane wave type supported by a Neveu-Schwarz two-form potential. This model is an open string version of the WZW model based on a non-semi-simple group previously studied by Nappi and Witten. The background we study is not conformally invariant. We consider a light-cone action for the sigma-model, compute the worldsheet propagator, and use it to exemplify a derivation of a time-dependent noncommutativity parameter.

¹ dolan@physics.unc.edu

² cnappi@princeton.edu

1. Introduction

Noncommutativity in string theory is a very interesting topic, as it may have important implications for the structure of spacetime. Noncommutativity has emerged in the context of open strings, starting from the treatment of open string field theory in [1]. More recently, it has reappeared in the context of Matrix theory compactified on a torus [2,3], and in the low energy description of strings in an electromagnetic background [4,5].

It is interesting to find other models in which noncommutativity emerges. In most of the examples currently known, the noncommutativity parameter is constant. An obvious task is to look for time-dependent noncommutativity parameters, especially given the recent interest in strings on time-dependent backgrounds [6-19].

In this paper we study an open string model, whose target space has a plane wave metric supported by a time-dependent Neveu-Schwarz two-form potential. This background was studied by Nappi and Witten [20] for closed strings. Here we are looking at an open string version, and by computing the worldsheet propagator we can derive a time-dependent noncommutativity parameter. It is important that the background is of the Neveu-Schwarz type: plane waves with Ramond fields remain commutative as the Ramond background amounts to the addition of a mass term to the action in light-cone gauge. In our case, for large values of the time parameter, our model reduces to a neutral string in a constant background B field [4,21], hence it is a good candidate for space-time noncommutativity.

In sect.2, we consider a light-cone action of the model. The mode expansion of a closed string version of this model has been explicitly exhibited in [22,23]. We compute the open mode expansion as a power series in a suitable parameter μ . This expansion is adequate to show noncommutativity. In sect.3 the worldsheet propagator is derived on the disk. In sect.4 we evaluate the propagator on the boundaries and compute a time-dependent noncommutativity parameter. The techniques used in this calculation are similar to those of [21] which analyzes strings in a $U(1) \times U(1)$ background.

2. A Time-Dependent Background

The Polyakov action coupling a string to a general metric and background Neveu-Schwarz field is

$$S = \int_{\Sigma} d\tau d\sigma [\sqrt{-\gamma} \gamma^{\alpha\beta} G_{MN} \partial_{\alpha} X^M \partial_{\beta} X^N + B_{MN} \epsilon^{\alpha\beta} \partial_{\alpha} X^M \partial_{\beta} X^N] \quad (2.1)$$

where we choose the string worldsheet Σ with Lorentz signature, and have rescaled the scalar worldsheet fields by $(2\sqrt{\pi\alpha'})^{-1}$ so that the X^M are dimensionless. We consider the time-dependent background provided by the Nappi Witten WZW model based on a non-semi-simple group, and adopt the same notation as in [20], with $X^M = (a_1, a_2, u, v)$, and u being identified with the time in the target space.

$$G_{MN} = \begin{pmatrix} 1 & 0 & \frac{a_2}{2} & 0 \\ 0 & 1 & -\frac{a_1}{2} & 0 \\ \frac{a_2}{2} & -\frac{a_1}{2} & b & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix}, \quad B_{MN} = \begin{pmatrix} 0 & u & 0 & 0 \\ -u & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}. \quad (2.2)$$

The Lorentz signature target space metric G_{MN} can be recognized as a plane wave metric [20]. The time-dependence is the u -dependence of B_{12} . Nappi and Witten checked that this model is exactly conformally invariant (*i.e.* to all orders in α') by showing the one-loop β function equations for the closed string backgrounds were satisfied, and then proving there were no higher order graphs.

In this paper, since we are interested in noncommutativity, we consider open string boundary conditions. Our case is not conformally invariant, and the background (2.2) satisfies the the Born-Infeld field equations* only for $N \neq u$

$$(D_M F_{NL})(1 - F^2)^{-1 LM} = 0 \quad (2.3)$$

where $(1 - F^2)^{-1 LM} = (1 + F)^{-1 LP} G_{PN} (1 - F)^{-1 NM}$ and $(1 - F)_{MN} \equiv G_{MN} - 2\pi\alpha' F_{MN}$. In our case $F_{MN} = B_{MN}$. For (2.2) the nonvanishing components of the Ricci tensor and affine connections are $R_{uu} = -\frac{1}{2}$, $\Gamma_{uj}^i = \frac{1}{2}\epsilon_{ij}^i$, $\Gamma_{ui}^v = -\frac{a_i}{4}$. It follows that $(D_M F_{NL})(1 - F^2)^{-1 LM} = \epsilon_{ij}(1 - F^2)^{-1 ju} = 0$, for $N \neq u$. But

$$(D_M F_{uL})(1 - F^2)^{-1 LM} = -\frac{u}{1 + u^2}. \quad (2.4)$$

As in [20], the sigma model action is (2.1) :

$$S = \int_{\Sigma} d\tau d\sigma [\sqrt{-\gamma} \gamma^{\alpha\beta} (\partial_{\alpha} a^i \partial_{\beta} a^i + 2\partial_{\alpha} u \partial_{\beta} v + b \partial_{\alpha} u \partial_{\beta} u + \epsilon_{ij} \partial_{\alpha} u \partial_{\beta} a^i a^j) + \epsilon^{\alpha\beta} \epsilon_{ij} u \partial_{\alpha} a^i \partial_{\beta} a^j]. \quad (2.5)$$

* We thank A. Hashimoto and K. Thomas [24] for pointing out an error in Eq.(2.3) in a previous version of this paper. A conformally invariant version of (2.2) is studied in [25], but its noncommutativity parameter although non-constant, is not time-dependent.

Although this action has a cubic interaction, if one treats it as a closed string theory, it is possible to find an exact mode expansion in the light-cone gauge [22,23]. However, in considering it as an open string theory, one has different boundary conditions which make the solution more complicated. Consequently, we will solve the theory only via a power series expansion. For simplicity, we work to lowest order in μ , where μ is a dimensionless constant, as this is sufficient to see noncommutativity. It is quite possible that another version of this model, differing from (2.5) via boundary terms, would lead to an exact mode expansion.

Although our background is not conformally invariant, we will consider a light-cone version of the sigma model in order to study open string propagators in a B-field with linear time dependence. We let

$$u = \mu\tau, \quad (2.6)$$

and write

$$S_{lc} = \int_{\Sigma} d\tau d\sigma [\eta^{\alpha\beta} \partial_{\alpha} a^i \partial_{\beta} a^i - 2\mu \partial_{\tau} v - b\mu^2 - \mu \epsilon_{ij} \partial_{\tau} a^i a^j + 2\epsilon_{ij} \mu\tau \partial_{\tau} a^i \partial_{\sigma} a^j]. \quad (2.7)$$

Here Σ is the string worldsheet with Minkowski metric $\eta^{\alpha\beta}$ with non-vanishing components $\eta^{\tau\tau} = -1, \eta^{\sigma\sigma} = 1$. We will use $\epsilon^{\tau\sigma} = 1$, and for the open string $-\infty \leq \tau \leq \infty, 0 \leq \sigma \leq \pi$.

With the fields a^i written in terms of $X \equiv a^1 + ia^2$ and $\tilde{X} \equiv a^1 - ia^2$, (2.7) becomes

$$S_{lc} = \int_{\Sigma} d\tau d\sigma [\eta^{\alpha\beta} \partial_{\alpha} X \partial_{\beta} \tilde{X} - 2\mu \partial_{\tau} v - b\mu^2 - \frac{i}{2}\mu(\partial_{\tau} X \tilde{X} - \partial_{\tau} \tilde{X} X) + i\epsilon^{\alpha\beta} \mu\tau \partial_{\alpha} X \partial_{\beta} \tilde{X}]. \quad (2.8)$$

The equations of motion and boundary conditions for the transverse fields are:

$$\begin{aligned} \square X - i\mu(\partial_{\sigma} X - \partial_{\tau} X) &= 0, & \square \tilde{X} + i\mu(\partial_{\sigma} \tilde{X} - \partial_{\tau} \tilde{X}) &= 0, \\ [\partial_{\sigma} X + i\mu\tau \partial_{\tau} X]|_{\sigma=0,\pi} &= 0, & [\partial_{\sigma} \tilde{X} - i\mu\tau \partial_{\tau} \tilde{X}]|_{\sigma=0,\pi} &= 0, \end{aligned} \quad (2.9)$$

where $\square \equiv -\partial_{\tau}^2 + \partial_{\sigma}^2 = 4z\bar{z}\partial_z\partial_{\bar{z}}$.

For large τ (so that τ can be considered constant), notice the similarity of the boundary condition in (2.9) with the boundary condition for an open string in a background B field. Since in the latter case the noncommutativity parameter is proportional to the background, this suggests we should expect here a noncommutativity parameter which depends on time.

The solution of (2.9) is given by the normal mode expansion for the transverse coordinates X and $\tilde{X}$, to first order in μ :

$$\begin{aligned}
X(\sigma, \tau) &= x_0 + a_0[\tau + \mu(-i\tau\sigma + \frac{i}{2}\tau^2)] \\
&\quad + \sum_{n \neq 0} a_n e^{-in\tau} [\frac{i}{n} \cos n\sigma + \mu((- \frac{1}{2n^2} - i\frac{\tau}{n}) \sin n\sigma + (\frac{i}{2n^2} + \frac{(\sigma - \tau)}{2n}) \cos n\sigma)] + O(\mu^2) \\
\tilde{X}(\sigma, \tau) &= \tilde{x}_0 + \tilde{a}_0[\tau - \mu(-i\tau\sigma + \frac{i}{2}\tau^2)] \\
&\quad + \sum_{n \neq 0} \tilde{a}_n e^{-in\tau} [\frac{i}{n} \cos n\sigma - \mu((- \frac{1}{2n^2} - i\frac{\tau}{n}) \sin n\sigma + (\frac{i}{2n^2} + \frac{(\sigma - \tau)}{2n}) \cos n\sigma)] + O(\mu^2)
\end{aligned} \tag{2.10}$$

We have derived (2.10) as follows. In (2.9) substitute $X(\sigma, \tau) = e^{i\frac{\mu}{2}(\tau+\sigma)} \phi(\sigma, \tau)$, and find

$$\begin{aligned}
\Box \phi &= 0, \\
[(\partial_\sigma + i\mu\tau\partial_\tau) \phi + i\frac{\mu}{2}(1 + i\mu\tau) \phi]_{\sigma=0, \pi} &= 0.
\end{aligned} \tag{2.11}$$

One such solution is $\phi(\sigma, \tau) = x_0 e^{-i\frac{\mu}{2}(\tau+\sigma)}$, corresponding to the constant mode $X(\sigma, \tau) = x_0$. A general solution to the wave equation $\Box \phi = 0$ is

$$\phi(\sigma, \tau) = f(\tau + \sigma) + g(\tau - \sigma). \tag{2.12}$$

So the constant solution above corresponds to $\phi(\sigma, \tau) = f(\tau + \sigma) = x_0 e^{-i\frac{\mu}{2}(\tau+\sigma)}$, and $g(\tau - \sigma) = 0$. To generate the solutions which provide the coefficients of a_0 and a_n in the normal mode expansion of $X(\sigma, \tau)$, we will try to find solutions $\phi(\sigma, \tau) = f(\tau + \sigma) + g(\tau - \sigma)$ satisfying the boundary conditions (2.11) via the power series expansions

$$\begin{aligned}
f(\tau + \sigma) &= \sum_{p=0}^{\infty} C_p (\tau + \sigma)^p \\
g(\tau - \sigma) &= \sum_{p=0}^{\infty} D_p (\tau - \sigma)^p
\end{aligned} \tag{2.13}$$

and

$$\begin{aligned}
f_n(\tau + \sigma) &= e^{-in(\tau+\sigma)} \sum_{p=0}^{\infty} C_p(n) (\tau + \sigma)^p \\
g_n(\tau - \sigma) &= e^{-in(\tau-\sigma)} \sum_{p=0}^{\infty} D_p(n) (\tau - \sigma)^p
\end{aligned} \tag{2.14}$$

respectively. A solution of (2.11) , in the form of (2.13) is

$$\begin{aligned}\mu\phi(\sigma, \tau) = & \mu\tau + \mu^2[-i\frac{3}{2}\tau\sigma] + \mu^3[\frac{1}{2}\tau^2\sigma + \frac{1}{6}\sigma^3 - \frac{9}{8}\tau\sigma^2 - \frac{3}{8}\tau^3 - \frac{\pi}{4}(\tau^2 + \sigma^2)] \\ & + i\mu^4[-\frac{1}{6}\tau^4 + \frac{21}{16}\tau^3\sigma - \tau^2\sigma^2 + \frac{21}{16}\tau\sigma^3 - \frac{1}{6}\sigma^4 + \pi(-\frac{3}{8}\tau^3 + \frac{5}{8}\tau^2\sigma - \frac{9}{8}\tau\sigma^2 + \frac{5}{24}\sigma^3) \\ & + \frac{\pi^2}{24}(\tau^2 + \sigma^2)] + O(\mu^5).\end{aligned}\tag{2.15}$$

where the functions f and g are given by

$$\begin{aligned}\mu f(\tau) = & \frac{\mu}{2}\tau - i\frac{3}{8}\mu^2\tau^2 - \mu^3\frac{\pi}{8}\tau^2 - \frac{5}{48}\mu^3\tau^3 + i\frac{31}{3 \cdot 128}\mu^4\tau^4 + i\mu^4(-\frac{\pi}{12}\tau^3 + \frac{\pi^2}{48}\tau^2) + O(\mu^5) \\ \mu g(\tau) = & \frac{\mu}{2}\tau + i\frac{3}{8}\mu^2\tau^2 - \mu^3\frac{\pi}{8}\tau^2 - \frac{13}{48}\mu^3\tau^3 - i\frac{95}{3 \cdot 128}\mu^4\tau^4 + i\mu^4(-\frac{7\pi}{24}\tau^3 + \frac{\pi^2}{48}\tau^2) + O(\mu^5).\end{aligned}\tag{2.16}$$

These expressions are derived iteratively, by considering the solution of (2.11) to some order μ^p , and then integrating the boundary condition to find the solution to order μ^{p+1} . Since finding a general form in arbitrary p , and summing these series to a closed form is difficult, we work to first order in μ . Note that although τ, σ could be rescaled to essentially eliminate μ , we keep it here to track the order in the power series solution of (2.11) . The series in (2.16) are reminiscent of hypergeometric functions. To derive the coefficient of a_n , we use the ansatz (2.14) to find

$$\phi_n(\sigma, \tau) = ie^{-in\tau} [\cos n\sigma + \mu((- \tau + \frac{i}{2n}) \sin n\sigma + (-i\sigma + \frac{1}{2n}) \cos n\sigma) + O(\mu^2)] \tag{2.17}$$

where $\phi_n(\sigma, \tau) = f_n(\tau + \sigma) + g_n(\tau - \sigma)$ with

$$\begin{aligned}f_n(\tau) = & ie^{-in\tau} [\frac{1}{2} + \mu(-\frac{i}{2}\tau) + O(\mu^2)] \\ g_n(\tau) = & ie^{-in\tau} [\frac{1}{2} + \mu(\frac{i}{2}\tau + \frac{1}{2n}) + O(\mu^2)].\end{aligned}\tag{2.18}$$

We then construct the normal mode expansion that satisfies (2.9) from

$$X(\sigma, \tau) = x_0 + e^{i\frac{\mu}{2}(\tau+\sigma)} a_0 \phi(\sigma, \tau) + e^{i\frac{\mu}{2}(\tau+\sigma)} \sum_{n \neq 0} a_n \phi_n(\sigma, \tau). \tag{2.19}$$

From (2.15) and (2.17) , we see that $X(\sigma, \tau)$ is given by an expansion where the coefficients of a_0, a_n are themselves a double power series in σ and τ . Although our open string model satisfies an equation of motion that can be simply related to the one-dimensional wave equation (2.9) , the particular boundary condition that is required substantially

complicates the form of the solution. (2.10) is reproduced by expanding (2.19) to first order in μ , using (2.15) and (2.17). Let $\mu \rightarrow -\mu$ to find $\tilde{X}(\sigma, \tau)$.

To quantize the theory in standard form, we reinsert the scale $2\sqrt{\pi\alpha'}$ so that $X, \tilde{X}$ become fields with length dimension, and find the canonical momenta:

$$\begin{aligned} P(\sigma, \tau) &= -\frac{\delta S}{\delta \partial_\tau X} = \frac{1}{4\pi\alpha'}(\partial_\tau \tilde{X} + i\frac{\mu}{2}\tilde{X} - i\mu\tau\partial_\sigma \tilde{X}) \\ \tilde{P}(\sigma, \tau) &= -\frac{\delta S}{\delta \partial_\tau \tilde{X}} = \frac{1}{4\pi\alpha'}(\partial_\tau X - i\frac{\mu}{2}X + i\mu\tau\partial_\sigma X). \end{aligned} \quad (2.20)$$

To first order in μ , we can invert the normal mode expansions in (2.10) as:

$$\begin{aligned} (1 + \frac{\mu}{2n})a_n &= \frac{1}{2\pi\sqrt{2\alpha'}} \int_0^\pi d\sigma \cos n\sigma [-in[X(\sigma, 0) + X(-\sigma, 0)] + [4\pi\alpha'[\tilde{P}(\sigma, 0) + \tilde{P}(-\sigma, 0)]]] \\ (1 - \frac{\mu}{2n})\tilde{a}_n &= \frac{1}{2\pi\sqrt{2\alpha'}} \int_0^\pi d\sigma \cos n\sigma [-in[\tilde{X}(\sigma, 0) + \tilde{X}(-\sigma, 0)] + [4\pi\alpha'[P(\sigma, 0) + P(-\sigma, 0)]]] \end{aligned} \quad (2.21)$$

for $n \neq 0$ and

$$\begin{aligned} x_0 &= \frac{1}{2\pi} \int_0^\pi d\sigma [X(\sigma, 0) + X(-\sigma, 0)]; \quad \tilde{x}_0 = \frac{1}{2\pi} \int_0^\pi d\sigma [\tilde{X}(\sigma, 0) + \tilde{X}(-\sigma, 0)] \\ \sqrt{2\alpha'}a_0 - i\frac{\mu}{2}x_0 &= 2\alpha' \int_0^\pi d\sigma [\tilde{P}(\sigma, 0) + \tilde{P}(-\sigma, 0)]; \quad \sqrt{2\alpha'}\tilde{a}_0 + i\frac{\mu}{2}\tilde{x}_0 = 2\alpha' \int_0^\pi d\sigma [P(\sigma, 0) + P(-\sigma, 0)]. \end{aligned} \quad (2.22)$$

The commutation relations which follow from canonical quantization $[X(\sigma, \tau), P(\sigma', \tau)] = i\delta(\sigma - \sigma')$, $[\tilde{X}(\sigma, \tau), \tilde{P}(\sigma', \tau)] = i\delta(\sigma - \sigma')$ are:

$$\begin{aligned} [a_m, \tilde{a}_n] &= 2(m - \mu)\delta_{m, -n}; \quad [a_m, a_n] = [\tilde{a}_m, \tilde{a}_n] = 0; \\ [x_0, \tilde{x}_0] &= 0; \quad [a_n, x_0] = [a_n, \tilde{x}_0] = [\tilde{a}_n, x_0] = [\tilde{a}_n, \tilde{x}_0] = 0 \text{ for } n \neq 0; \\ [x_0, \tilde{a}_0] &= i2\sqrt{2\alpha'} = [\tilde{x}_0, a_0]; \quad [x_0, a_0] = [\tilde{x}_0, \tilde{a}_0] = 0. \end{aligned} \quad (2.23)$$

3. The Propagator on the Disk

Having found a mode expansion, we compute the propagator, along the lines of [21]. In $z, \bar{z}$ coordinates (where z is in the upper half plane, since $0 \leq \sigma \leq \pi$), the equation of motion and boundary conditions for the propagator are:

$$\begin{aligned} 4z\bar{z}\partial_z\partial_{\bar{z}}X - 2\mu\bar{z}\partial_{\bar{z}}X &= 0, \quad 4z\bar{z}\partial_z\partial_{\bar{z}}\tilde{X} + 2\mu\bar{z}\partial_{\bar{z}}\tilde{X} = 0 \\ (\partial_z - \partial_{\bar{z}})X + \frac{\mu}{2}\ln z\bar{z}(\partial_z + \partial_{\bar{z}})X|_{z=\bar{z}} &= 0, \quad (\partial_z - \partial_{\bar{z}})\tilde{X} - \frac{\mu}{2}\ln z\bar{z}(\partial_z + \partial_{\bar{z}})\tilde{X}|_{z=\bar{z}} = 0 \\ 4\partial_z\partial_{\bar{z}}\langle X(z, \bar{z})\tilde{X}(\zeta, \bar{\zeta}) \rangle &= -2\mu z^{-1}\partial_{\bar{z}}\langle X(z, \bar{z})\tilde{X}(\zeta, \bar{\zeta}) \rangle = -2\pi\alpha'\delta^2(z - \zeta) \\ [(\partial_z - \partial_{\bar{z}})\langle X(z, \bar{z})\tilde{X}(\zeta, \bar{\zeta}) \rangle &+ \frac{\mu}{2}\ln z\bar{z}(\partial_z + \partial_{\bar{z}})\langle X(z, \bar{z})\tilde{X}(\zeta, \bar{\zeta}) \rangle]|_{z=\bar{z}} = 0. \end{aligned} \quad (3.1)$$

We will compute the propagator on the disk, and will use $z = e^{i(\tau+\sigma)}$, $\bar{z} = e^{i(\tau-\sigma)}$, $\zeta = e^{i(\tau'+\sigma')}$ and $\bar{\zeta} = e^{i(\tau'-\sigma')}$. In the above boundary conditions, the notation $|_{z=\bar{z}}$ denotes $z = |z|, \bar{z} = |z|$ at the $\sigma = 0$ endpoint and $z = |z|e^{i\pi}, \bar{z} = |z|e^{-i\pi}$ at $\sigma = \pi$. Assuming the commutation relations in (2.23), then for $|z| > |\zeta|$, the propagator to order μ is

$$\begin{aligned}
\langle X(z, \bar{z}) \tilde{X}(\zeta, \bar{\zeta}) \rangle &= \sqrt{2\alpha'} [a_0, \tilde{x}_0] (\tau + \mu(-i\tau\sigma + \frac{i}{2}\tau^2)) \\
&+ 2\alpha' \sum_{n=1}^{\infty} [a_n, \tilde{a}_m] e^{-in\tau} e^{-im\tau'} \left[-\frac{1}{nm} \cos n\sigma \cos m\sigma' \right. \\
&+ i\frac{\mu}{m} \cos m\sigma' \left(\left(-\frac{1}{2n^2} - \frac{i\tau}{n}\right) \sin n\sigma + \left(\frac{i}{2n^2} + \frac{(\sigma - \tau)}{2n}\right) \cos n\sigma \right) \\
&- i\frac{\mu}{n} \cos n\sigma \left(\left(-\frac{1}{2m^2} - \frac{i\tau'}{m}\right) \sin m\sigma' + \left(\frac{i}{2m^2} + \frac{(\sigma' - \tau')}{2m}\right) \cos m\sigma' \right) \Big] \\
&+ \mu(c_1\tau + c_0) \\
&= -i4\alpha' (\tau + \mu(-i\tau\sigma + \frac{i}{2}\tau^2)) \\
&+ 4\alpha' \sum_{n=1}^{\infty} e^{-in(\tau-\tau')} \left[\frac{1}{n} \cos n\sigma \cos n\sigma' \right. \\
&+ i\mu \cos n\sigma' \left(\left(\frac{1}{2n^2} + \frac{i\tau}{n}\right) \sin n\sigma - \left(\frac{i}{2n^2} + \frac{(\sigma - \tau)}{2n}\right) \cos n\sigma \right) \\
&- i\mu \cos n\sigma \left(\left(\frac{1}{2n^2} - \frac{i\tau'}{n}\right) \sin n\sigma' + \left(\frac{i}{2n^2} - \frac{(\sigma' - \tau')}{2n}\right) \cos n\sigma' \right) \\
&- \frac{\mu}{n^2} \cos n\sigma \cos n\sigma' \Big] \\
&+ \mu(c_1\tau + c_0).
\end{aligned} \tag{3.2}$$

We are free to add the function $\mu(c_1\tau + c_0)$ to the expression since it does not affect the equation of motion or the boundary condition for the propagator to first order in μ . For $|z| > |\zeta|$, the expression for $\langle \tilde{X}(z, \bar{z}) X(\zeta, \bar{\zeta}) \rangle$ is given by letting $\mu \rightarrow -\mu$ in the above propagator. In the $\mu \rightarrow 0$ limit, these propagators reduce to the open bosonic string propagator $\lim_{\mu \rightarrow 0} \langle X(z, \bar{z}) \tilde{X}(\zeta, \bar{\zeta}) \rangle = -2\alpha' (\ln |z - \zeta| + \ln |z - \bar{\zeta}|)$.

4. Time-Dependent Noncommutativity

To evaluate the noncommutativity parameter as defined from time ordering [26,4], we consider the propagator on the worldsheet boundary at $\sigma = 0$, then $z = |z| = e^{i\tau} \equiv \mathcal{T}$, and $\zeta = e^{i(\tau'+\sigma')} = |\zeta| = e^{i\tau'} = \mathcal{T}'$, so $\mathcal{T}, \mathcal{T}' > 0$. We will also consider the propagator at

$\sigma = \pi$, then $z = |z|e^{i\pi} = \mathcal{T}$ and $\zeta = |\zeta|e^{i\pi} = \mathcal{T}'$ so here $\mathcal{T}, \mathcal{T}' < 0$. Note that $\mathcal{T}$ is different from the worldsheet time τ .

$$\begin{aligned}
\langle X(z, \bar{z})\tilde{X}(\zeta, \bar{\zeta}) \rangle|_{\sigma=0} &= -i4\alpha' \left(\tau + \mu \frac{i}{2} \tau^2 \right) + \mu(c_1\tau + c_0) \\
&\quad - 4\alpha' \ln(1 - e^{-i(\tau-\tau')}) - 2\alpha' \mu i(\tau - \tau') \ln(1 - e^{-i(\tau-\tau')}) \\
&= -4\alpha' \ln(\mathcal{T} - \mathcal{T}') \\
&\quad + \mu(-2\alpha' \ln^2 \mathcal{T} - 2\alpha' \ln(\frac{\mathcal{T}}{\mathcal{T}'})) \ln(1 - \frac{\mathcal{T}'}{\mathcal{T}}) + (-c_1 i \ln \mathcal{T} + c_0) \\
\langle \tilde{X}(z, \bar{z})X(\zeta, \bar{\zeta}) \rangle|_{\sigma=0} &= -i4\alpha' \left(\tau - \mu \frac{i}{2} \tau^2 \right) - \mu(c_1\tau + c_0) \\
&\quad - 4\alpha' \ln(1 - e^{-i(\tau-\tau')}) + 2\alpha' \mu i(\tau - \tau') \ln(1 - e^{-i(\tau-\tau')})
\end{aligned} \tag{4.1}$$

Then at $\sigma = 0$:

$$\begin{aligned}
[X(\mathcal{T}), \tilde{X}(\mathcal{T})] &= T(X(\mathcal{T}) \tilde{X}(\mathcal{T}^-) - X(\mathcal{T})\tilde{X}(\mathcal{T}^+)) \\
&\equiv \lim_{\epsilon \rightarrow 0} (\langle X(\mathcal{T}) \tilde{X}(\mathcal{T} - \epsilon) \rangle - \langle \tilde{X}(\mathcal{T} + \epsilon) X(\mathcal{T}) \rangle), \quad (\text{for } \epsilon > 0) \\
&= \mu(-4i\alpha')(\pi \ln \mathcal{T} - i \ln^2 \mathcal{T}) \\
&= \mu 4\alpha' (\pi\tau + \tau^2) \equiv \Theta,
\end{aligned} \tag{4.2}$$

where we chose $c_1 = 2\pi\alpha'$, $c_0 = 0$, and use $\lim_{\epsilon \rightarrow 0}(\ln(1 + \epsilon) \ln \epsilon) = 0$. The noncommutativity parameter Θ is time-dependent.

At $\sigma = \pi$:

$$\begin{aligned}
\langle X(z, \bar{z})\tilde{X}(\zeta, \bar{\zeta}) \rangle|_{\sigma=\pi} &= -i4\alpha' \left(\tau + \mu(-i\tau\pi + \frac{i}{2}\tau^2) \right) + \mu(c_1\tau + c_0) \\
&\quad - 4\alpha' \ln(1 - e^{i(\tau'-\tau)}) \\
&\quad - 2\alpha' \mu i(\tau - \tau') \ln(1 - e^{-i(\tau-\tau')}) \\
\langle \tilde{X}(z, \bar{z})X(\zeta, \bar{\zeta}) \rangle|_{\sigma=\pi} &= -i4\alpha' \left(\tau - \mu(-i\tau\pi + \frac{i}{2}\tau^2) \right) + \mu(c_1\tau + c_0) \\
&\quad - 4\alpha' \ln(1 - e^{i(\tau'-\tau)}) \\
&\quad + 2\alpha' \mu i(\tau - \tau') \ln(1 - e^{-i(\tau-\tau')})
\end{aligned} \tag{4.3}$$

$$\begin{aligned}
[X(\mathcal{T}), \tilde{X}(\mathcal{T})] &= T(X(\mathcal{T}) \tilde{X}(\mathcal{T}^-) - X(\mathcal{T})\tilde{X}(\mathcal{T}^+)) \\
&\equiv \lim_{\epsilon \rightarrow 0} (\langle X(\mathcal{T}) \tilde{X}(\mathcal{T} + \epsilon) \rangle - \langle \tilde{X}(\mathcal{T} - \epsilon) X(\mathcal{T}) \rangle), \quad (\text{for } \epsilon > 0) \\
&= (-i4\alpha') \mu[-\pi \ln \mathcal{T} - i \ln^2 \mathcal{T}] \\
&= \mu 4\alpha' (-\pi\tau + \tau^2)
\end{aligned} \tag{4.4}$$

Thus for small μ , we have:

$$\begin{aligned}\Theta &= \mu \, 4\alpha' (\pi\tau + \tau^2) & \text{at } \sigma = 0, \\ \Theta &= \mu \, 4\alpha' (-\pi\tau + \tau^2) & \text{at } \sigma = \pi.\end{aligned}\tag{4.5}$$

For small τ , the theta parameter at the $\sigma = 0$ end of the string is minus that at the $\sigma = \pi$ end. This is the case for the neutral string in a constant background B field as well. In fact, although we have worked only to lowest order in μ , we can see directly from the equations of motion and boundary conditions (in $z, \bar{z}$) variables in (3.1), that in the limit of large z , *i.e.* large $i\tau$, a limit for which $z^{-1} \rightarrow 0$, that the system reduces to the neutral string with the identification $-\mu\tau = B$, a constant. (In the large τ limit, we note that $\ln|z|$ is approximately constant, in the sense that it is changing slowly, *i.e.* its derivative $|z|^{-1}$ is small. Therefore, for large τ the noncommutativity parameter becomes constant, and our model is similar to the neutral string.) For large τ , using the neutral string expressions, we find the noncommutativity parameter be time-dependent:

$$\begin{aligned}\Theta &= -4\alpha' \pi B = 4\alpha' \mu \pi \tau & \text{at } \sigma = 0, \\ \Theta &= 4\alpha' \pi B = -4\alpha' \mu \pi \tau & \text{at } \sigma = \pi.\end{aligned}\tag{4.6}$$

We have shown that our model exhibits noncommutativity for both small and large τ . The expectation is that the model will remain noncommutative with a time-dependent noncommutativity parameter for all times.

Acknowledgements

It is a pleasure to thank Edward Witten for discussions. LD is grateful to Princeton University and the Institute for Advanced Study for their hospitality during the summer 2002, and to the Aspen Center for Physics. She was supported in part by U.S. Department of Energy, Grant No. DE-FG 05-85ER40219/Task A. CRN is supported in part by NSF grant PHY-0140311. [Any opinion, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the National Science Foundation.]

References

- [1] E. Witten, “Noncommutative Geometry and String Field Theory,” Nucl. Phys. **B268** (1986) 253.
- [2] A. Connes, M.R. Douglas, and A. Schwarz, “Noncommutative Geometry and Matrix Theory: Compactification on Tori,” JHEP 9802 (1998) 003, hep-th/9711162.
- [3] M. Douglas and N. Nekrasov, “Noncommutative Field Theory”, Rev. Mod. Phys. **73** (2001) 977, hep-th/0106048.
- [4] N. Seiberg and E. Witten, “String Theory and Noncommutative Geometry”, JHEP **9909** (1999) 032, hep-th/9908142.
- [5] A. Abouelsaood, C. Callan, C. Nappi, and S. Yost, “Open Strings in Background Gauge Fields” Nuclear Physics **B280** (1987) 599.
- [6] G. Horowitz and A. Stief, “Strings in Strong Gravitational Fields”, Physical Review **D42** (1990) 1950.
- [7] A. Sen, “Time Evolution in Open String Theory”, hep-th/0207105.
- [8] S. Hemming, E. Keski-Vakkuri, P. Kraus, “Strings in the Extended BTZ Space-Time”, hep-th/0208003.
- [9] L. Cornalba and M. Costa, “A New Cosmological Scenario in String Theory”, Phys. Rev. **D66** (2002) 066001, hep-th/0203031.
- [10] J. Figueroa-O’Farrill and Joan Simon, “Generalised Supersymmetric Fluxbranes”, JHEP 0112 (2001) 011, hep-th/0110170.
- [11] A. Hashimoto, S. Sethi, “Holography and String Dynamics in Time Dependent Backgrounds”, hep-th/0208126.
- [12] M. Alishahiha and S. Parvizi, “Branes in Time-Dependent Backgrounds and AdS/CFT Correspondence”, hep-th/0208187.
- [13] H. Liu, G. Moore, and N. Seiberg, “Strings in Time Dependent Orbifolds”, hep-th/0204168, hep-th/0206182.
- [14] G. Horowitz and J. Polchinski, “Instability of Space-Like and Null Orbifold Singularities”, hep-th/0206228.
- [15] E. Gimon, L. Pando Zayas and J. Sonnenschein, “Penrose Limits and RG Flows”, hep-th/0206033.
- [16] H. Lewis and W. Riesenfeld, “An Exact Quantum Theory of the Time-Dependent Harmonic Oscillator and of a Charged Particle in a Time-Dependent Electromagnetic Field”, Jour. Math. Phys. **10** (1969) 1458.
- [17] D. Khandekar and S. Lawande, “Exact Propagator for a Time-Dependent Harmonic Oscillator with and without a Singular Perturbation”, Jour. Math. Phys. **16** (1975) 384.
- [18] C. Duval, Z. Horvath, and P. Horvathy, “Vanishing of the Conformal Anomaly for Strings in a Gravitational Wave”, Phys. Lett. **B313** (1993) 10.

- [19] C. Duval, Z. Horvath, and P. Horvathy, “Strings in Plane-fronted Gravitational Waves”, *Mod. Phys. Lett.* **A8** (1993) 39.
- [20] C. Nappi and E. Witten, “A WZW Model Based on a Non-semi-simple Group”, *Phys. Rev. Lett.* **71** (1993) 3751; hep-th/9310112.
- [21] L. Dolan and C. Nappi, “A Scaling Limit With Many Noncommutativity Parameters”, *Phys. Lett.* **B504** (2001) 329; hep-th/0009225.
- [22] P. Forgacs, P. Horvathy, Z. Horvath, and L. Palla, “The Nappi-Witten string in the light-cone gauge”, hep-th/9503222.
- [23] J.G. Russo and A.A. Tseytlin, “Constant Magnetic Field in Closed String Theory: An Exactly Solvable Model”, *Nucl. Phys.* **B448** (1995) 293, hep-th/9411099.
- [24] A. Hashimoto and K. Thomas, “Dualities, Twists, and Gauge Theories with Non-Constant Non-Commutativity”, hep-th/0410123.
- [25] S. Stanciu and A. Tseytlin, “D-branes in Curved Spacetime: Nappi-Witten Background,” *JHEP* **9806**, 010 (1998) hep-th/9805006.
- [26] V. Schomerus, “D-Branes and Deformation Quantization,” *JHEP* **9906** (1999) 030; hep-th/9903205.